



Hamilton County Public Health - Epidemiology and Assessment

Chlamydia and Gonorrhea Quarterly Report

Kevin Strobino, MPH, Epidemiologist

Chlamydia Infections by Month (2015-2016)

Table 1. Chlamydia Cases by Month for Hamilton County Residents


Month	Cases of Chlamydia 2015	Cases of Chlamydia 2016
January	627	563
February	504	554
March	611	634
April	612	604
May	560	566
June	596	570
July	613	
August	663	
September	632	
October	548	
November	522	
December	582	
Total	7,070	3,491

This report was created as a surveillance effort to help prevent new cases of chlamydia and gonorrhea within Hamilton County. Table 1 displays the total number of chlamydia cases for Hamilton County residents (at diagnosis) from 2015 and the second quarter of 2016 on a monthly basis. Only chlamydia cases that have been reported to the CDC were counted for analysis purposes in this report. In 2015, the highest number of chlamydia cases was reported in August (663 cases). Through Q2 of 2016, the highest number of chlamydia cases occurred in March (634 cases). There have been 581.8 chlamydia cases per month through Q2 2016, about 1% lower than the monthly average of 589.2 in 2015. Assuming negligible variance in cases reported between months, 2016 is presently on pace to have 6,982 reported chlamydia cases, an estimated decrease from 2015 of 88 cases.

Chlamydia cases are derived from data in the Ohio Disease Reporting System and represent only those cases reported to the CDC. These data are provisional and subject to change when additional data are reported. Cases are selected based on address at diagnosis. Source: Ohio Department of Health (ODH), STD Surveillance. Data reported as of 7/25/2016.

Surveillance of Chlamydia Cases in Hamilton County

One way to monitor chlamydia infections within Hamilton County is through the use of surveillance control charts. Factors that these control charts show are the number of chlamydia cases for each month (black diamonds), control limits (red or green dashed lines), and the average number of cases (solid blue line). Control charts are used to detect unexpected events, such as a single point outside of the control limit, consecutive points above or below the average line, or two or three consecutive points near a control limit. When anomalies such as these occur it may be beneficial to examine events surrounding the anomalies in order to devise a strategy to reduce the number of cases in subsequent months or see which strategies already in place are working. Figure 1 shows the control chart for chlamydia infections from July 2014 to June 2016. All of the months in this time-frame fell within the control limits for the monthly number of infections. The average number of cases was calculated from August 2013 to July 2014 (568.6).



Chlamydia cases are derived from data in the Ohio Disease Reporting System and represent only those cases reported to the CDC. These data are provisional and subject to change when additional information is reported. Cases are selected based on address at diagnosis. Source: Ohio Department of Health, STD Surveillance. Data reported as of 7/25/2016.




Demographics with High Risk for Chlamydia Infection

Certain demographic groups are at higher risk for chlamydia infection. Table 2 shows the percentage of chlamydia cases from 2015 and 2016 based on race, age and sex. Nearly 75 percent of the chlamydia cases from 2015 occurred among black Hamilton County residents; that figure is lowering towards 72% in 2016. About 67% of chlamydia cases were between the ages of 15-24, and the majority of diagnosed cases in 2015 and 2016 were among female Hamilton County residents. Identifying these at-risk groups allows public health and health care the opportunity to create focused intervention methods for preventing the spread of chlamydia. Figure 2 further classifies the differences among race/sex groups from 2013 to Q2 2016. The demographics from 2013 to 2016 are similar, with a more equitable distribution of cases emerging as the largest group, black females, continues to make up a lower percentage of all chlamydia cases (50.2% through Q2 2016).

Table 2. Demographics of Chlamydia Cases

	2015		2016	
	#	%	#	%
Race				
Black	3,376	74.7	1,757	71.6
White	936	20.7	565	23.0
Other	215	4.8	133	5.4
Age				
<1	1	0.0	0	0.0
1-14	116	1.6	47	1.4
15-24	4,843	68.6	2,348	67.3
25-34	1,636	23.2	848	24.3
35-44	339	4.8	188	5.4
45-54	95	1.4	39	1.1
55-64	27	0.4	16	0.5
>65	5	0.1	3	0.1
Sex				
Male	2,073	29.3	1,055	30.2
Female	4,997	70.7	2,436	69.8

Figure 2. Race/Sex Distribution of Chlamydia Cases, Hamilton County



These data are provisional and subject to change when additional data are reported. Chlamydia cases between January 2015 and June 2016 were used for analysis. Cases were selected based on address at diagnosis. Source: Ohio Department of Health, STD Surveillance. Data reported as of 7/25/2016. Percentages may not total to 100 percent due to rounding. Percentages for demographics are based only on cases that had valid information within the required fields.

Gonorrhea Infections by Month (2015-2016)

Table 3. Gonorrhea Cases by Month for Hamilton County Residents

Month	Cases of Gonorrhea 2015	Cases of Gonorrhea 2016
January	265	240
February	179	216
March	233	248
April	229	224
May	212	220
June	237	242
July	231	
August	239	
September	244	
October	258	
November	223	
December	247	
Total	2,797	1,390


Table 3 displays the total number of gonorrhea cases for Hamilton County residents (at diagnosis) from 2015 and the second quarter of 2016 on a monthly basis. Only gonorrhea cases that have been reported to the CDC were counted for analysis purposes in this report. In 2015, the highest number of gonorrhea cases was reported for January (265 cases). Through Q2 of 2016, the highest number of gonorrhea cases occurred in March (248 cases). The average number of gonorrhea cases per month was 233.1 and 231.7 for 2015 and through Q2 2016, respectively. Assuming negligible variance in cases reported between months, 2016 is presently on pace to have 2,780 reported gonorrhea cases, which would be a small decrease of 17 cases from 2015.

Gonorrhea cases are derived from data in the Ohio Disease Reporting System and represent only those cases reported to the CDC. These data are provisional and subject to change when additional data are reported. Cases are selected based on address at diagnosis. Source: Ohio Department of Health (ODH), STD Surveillance. Data reported as of 7/25/2016.



Surveillance of Gonorrhea Cases in Hamilton County

One way to monitor gonorrhea infections within Hamilton County is through the use of surveillance control charts. Factors that these control charts show are the number of gonorrhea cases for each month (black diamonds), control limits (red or green dashed lines), and the average number of cases (solid blue line). Control charts are used to detect unexpected events, such as a single point outside of the control limit, consecutive points above or below the average line, or two or three consecutive points near a control limit. When anomalies such as these occur it may be beneficial to examine events surrounding the anomalies in order to devise a strategy to reduce the number of cases in subsequent months or see which strategies already in place are working. Figure 3 illustrates the control chart for gonorrhea infections from July 2014 to June 2016. All of the months within this time frame fell below the upper control limit for number of gonorrhea infections. The average number of cases was calculated from October 2014 to September 2015 (224.7).




Gonorrhea cases are derived from data in the Ohio Disease Reporting System and represent only those cases reported to the CDC. These data are provisional and subject to change when additional information is reported. Cases are selected based on address at diagnosis. Source: Ohio Department of Health, STD Surveillance. Data reported as of 7/25/2016.

Demographics with High Risk for Gonorrhea Infection

Individuals with certain demographics are more likely to be infected with gonorrhea. Table 4 shows the percentage of gonorrhea cases from 2015 and 2016 based on race, age and sex. About 80 percent of the gonorrhea cases from 2015 and through Q2 2016 occurred among black Hamilton County residents. Over half of gonorrhea cases were between the ages of 15-24. Identifying these aforementioned at-risk groups allows public health and health care the opportunity to create focused intervention methods for preventing the spread of gonorrhea. Figure 4 further classifies the differences among race/sex groups from 2013 to 2016. There has been a large reduction in the percentage of cases that are in the majority group, black females, while the percentage of cases that are black males and white females have steadily increased.

Table 4. Demographics of Gonorrhea Cases

	2015		2016	
	#	%	#	%
Race				
Black	1,633	80.6	815	78.4
White	336	16.6	181	17.4
Other	58	2.9	43	4.1
Age				
<1	0	0	0	0
1-14	23	0.8	11	0.8
15-24	1,568	56.1	723	52.1
25-34	814	29.1	433	31.2
35-44	233	8.3	132	9.5
45-54	101	3.6	62	4.5
55-64	45	1.6	23	1.7
>65	11	0.4	4	0.3
Sex				
Male	1,266	45.3	650	46.8
Female	1,531	54.7	740	53.2



These data are provisional and subject to change when additional data are reported. Gonorrhea cases between January 2015 and June 2016 were used for analysis. Cases were selected based on address at diagnosis. Source: Ohio Department of Health, STD Surveillance. Data reported as of 7/25/2016. Percentages may not total to 100 percent due to rounding. Percentages for demographics are based only on cases that had valid information within the required fields.