



Hamilton County Public Health - Epidemiology and Assessment

Syphilis Quarterly Report

David Carlson, MPH, Epidemiologist

Syphilis Infections by Month, Hamilton County, Ohio (January 2012 – December 2013)

Table 1. Syphilis Cases by Month for Hamilton County Residents


Month	Cases of Syphilis 2012	Cases of Syphilis 2013
January	20	30
February	31	35
March	37	33
April	51	30
May	41	28
June	29	23
July	31	35
August	40	27
September	44	32
October	32	34
November	37	17
December	38	20
Total	431	344

This report was created as a surveillance effort to help prevent new cases of syphilis within Hamilton County. Table 1 displays the breakdown of syphilis cases for Hamilton County residents from 2012 and 2013 on a monthly basis. Only syphilis cases that have been reported to the CDC were counted for analysis purposes in this report. In 2012, the highest number of syphilis cases occurred in April (51 cases). In 2013, the highest number of syphilis cases occurred in February and July (35 cases). The average number of syphilis cases per month were 35.9 and 28.7 for the years 2012 and 2013, respectively. In the fourth quarter of 2013, there were 36 fewer cases of syphilis compared to the fourth quarter of 2012. Subsequent reports will allow for a better comparison of 2013 cases as data are subject to change as more information is acquired.

Syphilis cases are derived from partner services data in the Ohio Disease Reporting System and represent only those cases reported to the CDC. These data are provisional and subject to change when additional data are reported. Cases are selected based on address at diagnosis. Source: Ohio Department of Health (ODH), STD Surveillance. Data reported as of 2/2/2014.

Surveillance of Syphilis Cases Diagnosed in Hamilton County, Ohio (2012 - 2013)

One way to monitor syphilis infections within Hamilton County is through the use of surveillance control charts. Factors that these control charts show are the number of syphilis cases for each month (black diamonds), control limits (red or green dashed lines), and the average number of cases (solid blue line). Control charts are used to detect unexpected events, such as a single point outside of the control limit, consecutive points above or below the average line, or two to three consecutive points near a control limit. When anomalies such as these occur it may be beneficial to examine events surrounding the anomalies in order to devise a strategy to reduce the number of cases in subsequent months or to see which strategies already in place are effective. Figure 1 illustrates the control chart for syphilis infections from January 2012 to December 2013. All of the months in this time frame fell below the upper control limit for number of syphilis infections. Future control charts will give a better understanding of the case counts for 2013. The monthly average number of cases (33.5) was calculated from January 2010-December 2012.



Syphilis cases are derived from partner services data in the Ohio Disease Reporting System and represent only those cases reported to the CDC. These data are provisional and subject to change when additional information is reported. Cases are selected based on address at diagnosis. Source: ODH, STD Surveillance. Data reported as of 2/2/2014.




Demographics and Social Factors with High Risk for Syphilis Infection

Individuals with certain demographics and social factors are more likely to become infected with syphilis. Table 2 shows the percentage of syphilis cases from 2012 and 2013 based on race, age, sex, and additional demographics. Approximately 75 percent of the syphilis cases from 2012 and 2013 occurred among black Hamilton County residents. Nearly 40 percent of the syphilis cases in 2012 were between the ages of 15-24. A switch in demographics occurred in January - December 2013 as a higher percentage of cases were between the ages of 25-34 and a higher percentage of cases were male, compared to 2012. Identifying these aforementioned at-risk groups allows public health and health care the opportunity to create specific intervention methods for preventing the spread of syphilis. Figure 2 further classifies the differences in race/sex groups from 2012 to 2013. The largest changes occurred among the percentage of black male, white male, and black female cases. The demographics for 2013 are subject to change as more cases are identified in 2013.

Table 2. Demographics of Syphilis Cases

	Jan. - Dec. 2012		Jan. - Dec. 2013	
	#	%	#	%
Race				
Black	344	79.8	247	71.8
White	57	13.2	73	21.2
Other	30	7.0	24	7.0
Age				
<1	6	1.4	7	2.0
1-14	1	0.2	1	0.3
15-24	165	38.3	104	30.2
25-34	128	29.7	120	34.9
35-44	62	14.4	69	20.1
45-54	54	12.5	36	10.5
55-64	13	3.0	7	2.0
>65	2	0.5	0	0.0
Sex				
Male	205	47.6	215	62.5
Female	226	52.4	129	37.5
Behavior*				
MSM	63 of 159	39.6	83 of 192	43.2
HRHF	121 of 223	54.3	62 of 118	52.5



These data are provisional and subject to change when additional data are reported. Syphilis cases between January 2012 and December 2013 were used for analysis. Cases were selected based on address at diagnosis. Source: ODH, STD Surveillance. Data reported as of 2/2/2014. Percentages may not total to 100 percent due to rounding. *Cases were missing information from fields used to determine transmission. Percentages for behavior are sex-specific and based only on cases that had valid information within the required fields. High risk heterosexual females (HRHF) are women who self-identified as participating in sex with a known MSM, HIV+, IDU, or anonymous person. HRHF status is also determined from factors such as having sex while intoxicated, exchanging sex for drugs, or having previous STIs.

Stages of Syphilis Infection: Hamilton County

Syphilis infections are organized into different stages based on the clinical presentation of disease and duration of infection. Congenital syphilis cases are cases of syphilis in which the infection is transferred from mother to infant during pregnancy or delivery. Congenital syphilis cases serve as key indicators of community health as this stage of infection is easily preventable when proper healthcare is present. Transmission of syphilis is possible during primary, secondary, and early latent stages of disease. In particular, primary and secondary infections are considered highly infectious stages. During late latent syphilis the patient is no longer infectious and has no symptoms; however if the patient does not receive treatment the disease can develop into neurological problems, possibly leading to death. As seen in Figure 3, the majority of the monthly cases in 2012 and 2013 are diagnosed as late latent cases.

